





IMPORTANT SAFETY INFORMATION FOR PATIENTS RECEIVING TREATMENT WITH **TECVAYLI or TALVEY**

Carry this card with you at all times. SHOW THIS CARD to any healthcare professional involved in your care and if you go to the emergency room.

TECVAYLI[®] and TALVEY[™] REMS PATIENT WALLET CARD

FOR HEALTHCARE PROFESSIONALS

may be tatal or lite-threatening. CR5 may involve multiple organ systems. including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) which TALVEY therapy can cause cytokine release syndrome (CRS) or neurologic toxicity, bn6 IJYAVDET : WONY GUOHZ UOY NOITAMAOHI YTERAS TNATAOMI

Complete the Patient Wallet Card based on the drug prescribed and provide the Patient Wallet Card to the patient.

This Patient has received: 🗆 TECVAYLI 🗖 TALVEY (Please check one.)

First Treatment Dose:

Office Phone Number:

Bi-Weekly Dosing (Please check one.) **Dosing** :elubedo2 gnisod

Dates of Injections:

:z əsog dn-dəiş • :Lesou qu-qest

Palies only if patient is receiving TALVEY at a treatment dose of 0.8 mg/kg

.¿cosog dn-doig .

Healthcare Setting Name:

:19dmuN 9nod9 21uoH 1931A sigoloonO gnitearT to ameN

FOR THE PATIENT

Call your healthcare professional or get emergency help right away if you recognize any of these symptoms:		
Symptoms of Cytokine Release Syndrome (CRS): • Fever (100.4°F or higher) • Difficulty breathing • Chills • Fast heartbeat • Dizziness or light-headedness • Feeling anxious • Confusion or restlessness • Headache	Symptoms of neurologic • Headache • Feeling confused or disoriented • Being less alert or aware • Feeling sleepy • Low energy (lethargy) • Slowness or difficulty thinking • Seizures • Memory Loss • Jerking movements • Rigid muscles • Feeling restless	 problems: Numbness and tingling (feeling like "pins and needles") Trouble speaking or writing Muscle spasms Shaking (tremors) Problems walking Hearing Loss Muscle weakness in your body or face Double vision Burning, throbbing or stabbing pain

You should always ask your doctor about taking other medications while taking TECVAYLI or TALVEY.



IMPORTANT TO REMEMBER: You may be asked to stay in the hospital for 48 hours after administration of all doses within the step-up dosing schedule. If you have <u>any</u> of these symptoms, call your doctor or seek emergency medical attention right away! These are not all of the possible symptoms of **TECVAYLI** or **TALVEY**. Tell your doctor if you have any symptom that bothers you or does not go away.